Standing Medicine People

A guide to common medicines found in south central Manitoba

Ji-mikigaadegin mashkikiin omaa zhaawanong Manitoba akiing



In partnership with Long Plain First Nation, Swan Lake First Nation, Rolling River First Nation, and Food Matters Manitoba

Owe wiidanokiindiwag Ginooshkodeyaang, Gaabiskigamaag, Ditibineyaziibi dago Food Matters Manitoba Standing Medicine People refers to those medicines found on the land, sweet flag, berries, and other plants used to bring balance to our bodies. Knowledge about the standing medicine people is limited, changes have taken place on our lands that have greatly impacted our use and understanding of this resource.

Among those changes are impacts to the environment, including pollution, development, and agricultural impacts. There are also societal impacts that have impacted our awareness of the medicines harvested from the land.

The vision of this project is to identify safe places to harvest medicinal plants with a reduced risk of adverse health effects from environmental contamination. With safe harvesting locations and baseline information First Nation communities will be enabled to once again share, use and promote Anishinabe Knowledge and practices that lead to more balanced and healthy lifestyle for many community members.

This guide book is meant to assist community members interested in harvesting berries and plants for personal use. There are brief descriptions of the plants and habitats, as well as general information on how to harvest. For further information about the medicine of these plants, offer your tobacco to a knowledgeable person and commit to learn.

We would like to acknowledge David Daniels, Carol Beaulieu, Roberta Morrissette and Elvin Huntinghawk for their support and the people of Long Plain, Swan Lake, and Rolling River for their contributions to the project.

Miigwech,

Daniel Gladu Kanu, Tim Stevenson, David Heinrichs, Nicole Lamy, and Christina Janzen

Food Matters Manitoba Standing Medicine People Project Staff

Gaa-niibawiwaad Mashkikiin ijigaadewan ini mashkikiin gaamikigaadegin akiing. Wiike, mawinzwaan dago bebakaan netaawigigin gaa-minoshkaagowang giiyawinaaning. Bangii eta gikenjigaadewan ini mashkikiin. Bebakaan gegoon gaagii-bi-izhisegin gidakiiminanan gii-meshkwajise aaniin ezhigikendamang ono mashkikiin aaniin gaye ji-aabajichigaadegin.

Iwe aaniin ezhi-meshkwajiseg bagwajaya'ii, daabishkoo bichiboojigaadeg aki, onachigewinan dago gitigewinan. Zhigwa miinawaa aaniin ezhiseg owe ji-gikendamang ji-ondinamang mashkiki akiing onji.

Owe izhichigewin ji-wiiji'iwemagak aandi ji-ondinigaadegin mashkikiin gii-bichiboojigaadesinogin bebakaan gegoon gaa-izhichigaadeg aki. Gikenjigaadeg aandi geonizhishingiban ji-ondinigaadegiban mashkikiin, odaagashkitoonaawaa' Anishinaabeg ji-waawiindamaadiwaad aandi ge-ondinamawaapan mashkikiin, aaniin gaye geinaabajitoowaapan, nawach daa-mamino-ayaawag.

Owe mazina'igan ji-wiiji'igowaad Anishinaabeg aandi ji-mawinzowaad, aandi gaye ji-ondinamowad gegoon netaawiginig. Wawozhibii'igaadewan aaniin ayizhinaagwakin netaawigigin, aandi gaye ge-mikigaadegibaniin. Giiyaabi gegoon noonde-gikendaman ono mashkikiin, miizh asemaa awiya gekendang, mii imaa onji gikendamaan.

Ninaakomaanaanig ingi onji Ginooshkodeyaang, Gaabiskigamaag, Ditibineyaziibiing gaa-gii-wiiji'iwewaad owe dazhiikigaadeg.

Miigwech

Highbush Cranberry (Viburnum opulus)

- Plant grows to 4 meters tall
- Opposite branching
- Deciduous shrub

Leaves similar in shape to maple leaf

Flowers in white clusters

Where to Harvest

Found near rivers and streams. Often around aspen in woods or bush where it has access to sunlight.

How to Harvest

Harvest by pulling the fruit off the branch. Clean and share your harvest.

When to Harvest

Harvest in October, when you can smell cranberry and the fruit is bright red.

Uses

Can be used for sauces and jams. Tastes very tart, sweeter after first frost.

To find out more about other types of uses from the cranberry medicinal plant, offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.



Aniibimin(an)

- Netaawiging 4 (m) diba'iganaatig apiitakidewan
- Edawaya'ii odikwaniwan
- Giitwaami-aniibiishiwan
- Aniibiishan daabishkoo ziizibaakwadwaatig
- Waabishki-waabiganiwan

Endazhiging

Mikigaade jiigi-ziibiing dago ziibiinsing. Bagwajaya'ii azaadiikaang, aandi igo ge-zaagaatenig.

Gidaa-mawinz odikwaning onji. Biini' ge-izhi-maamiigiweyan. Gidaagiizizekwaadaan gemaa baashkiminisigaage. Ziiwiskipogwad, nawach zhiiwan ishkwaa nitam mashkwaakwading.

October mawinzon apii minaandaman aniibiminan dago aakomiskwaagin.

Giiyaabi qegoon noonde-qikendaman aaniin qe-danaabajitooyamban aniibiminan, miizh asemaa gichi-anishinaabeg gemaa awiya gekendang gechiwaag aki.





Saskatoon (Amelanchier alnifolia)

- Plants grows 1 to 5 metres tall
- Alternate branching
- Loses leaves in winter (Deciduous)
- Leaves are oval to circular in shape
- Flowers white in groups of 3 to 20, each flower with 5 white petals

Where to Harvest

Found in open areas where the plants have access to sunlight, but can also be found in forests. In moist but well-drained areas. Look for hills.

How to Harvest

Harvest by pulling the fruit off the branches. Fruit is blue-purple when ripe and round. Clean and share your harvest.

When to Harvest

Harvest in summer, July.

Uses

Fruit can be eaten raw, made into a jelly, or frozen for later use.

To find out more about other types of uses from the Saskatoon medicinal plant offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.







Ozigwaakomin(ag)

- Bezhig (1) ako Naanan (5) apiitakidewan
- Bebakaan wawodikwaniwan
- Binibagaa apii biboong
- Zhaagawaawan gemaa waawiyaawan aniibiishan.
- Niswi (3) ako (20) ako waabigwaniwan, naananobaganoon (5) waabigwaniin

Endazhiging

Mishawaya'ii nitaawigin aandi ezhi-zaagaateg, megwaakwaang dash gaye daa-mikigaadewan aandi dipaabaaweg nitaa-iskising dash. Andone'an wajiin.

Mawinzon odikwaning onji. Ozhaawashko-apisiziwag apii adisowaad, waawiyeziwag gaye.

Biini', maamiigiwen. Gidaa-ashkamaag, gidaa-baashkiminisigaageg gemaa daa-mashkawaakwajimaawag naagach onji.

Niibing mawinzon, July aginzod.

Giiyaabi noonde-qikendaman aaniin qe-danaabajiwindwaaban ozigwaakominag, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.

Chokecherries (Prunus virginiana)

Plant grows 1 to 6 metres tall

Deciduous shrub with alternate branching. Branches often covered in a black fungus known as Black Knot

Leaves oval but coming to a point at the tip. Bright green on top

Flowers in clusters. Petals white, Each flower with 5 petals

Where to Harvest

Spread throughout open areas and in woods or bush.

How to Harvest

Harvest by carefully pulling the fruit from the plant. Fruits, red drupelets, in dense clusters, about 1 cm across, that fall intact.

When to Harvest

Harvest in summer, late July.

Uses

Can be eaten fresh, or crushed and dried to be used later in cooking. Chokecherry can also be used to make jams and sauces.

To find out more about other types of uses from the Chokecherrry medicinal plant offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.



Azasowemin(an)

- Bezhig (1) ako Ningodwaaswi (6) diba'iganaatig apiitakidewan.
- Giitwaami-aniibiishiwan, bebakaan odikwaniwan. Naanigoding makade-wazhashkwedowan odikwanan
- Zhaagawaawan aniibiishan, jaajiibwaawan onakwaya'ii, aakoashkaabaganoon gaye imaa
- Okoginoon waabigwaniin. Waabishkaawan. Naanobagananoon endasingin

Endazhiging

Mishawaya'ii nitaawigin abooshke megwaakwaang gemaa noopimiing.

Mawinzon odikwaning onji. Aako-miskwaawan gemaa makadewaawan apii atisingin. Nawach igo ziiwiskaawan.

Niibing mawinzon, ani-giizhaginzod July.

Giiyaabi noonde-qikendaman aaniin qe-danaabajiwindibaniin azasoweminan, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.





Wild Raspberry (Rubus idaeus)

- Erect, perennial shrub, 1-2 m tall stems (canes) upright, biennieal, prickly, often with gland tipped hairs, bark shredding, yellow to cinnamon brown
- Leaves, compound, 3-5 leaflets per leaf, egg shaped, sharply pointed double saw toothed.
- Flowers, single or small clusters, drooping white, 8-12 cm

Where to Harvest

Found in thickets, clearings, and open woods.

How to Harvest

Harvest by carefully pulling the fruit from the plant. Fruits, red drupelets, in dense clusters, about 1 cm across, that fall intact.

When to Harvest

Harvest in summer months.

Uses

Can be eaten raw or canned for later use.

To find out more about other types of uses from the wild red raspberry medicinal plant offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.



Miskomin(ag)

- Badakidewan, endasowaki nitaawigin, bezhig (1) ako niizh (2) apiitakidewan, neniizhwaaki gaye, gaashaawan, ningoding onakwaya'ii dibaabagizi, biniwanagekizi, waabishkizaawizi gemaa ozaawizi
- Aniibiishan okoginoon, niswi (3) ako naanobaganoon (5) dasobaganoonsiwanoon, zhaagawaawan, aako-jiibo-gidagaawan
- Waabiwaniwan, bezhigobagaawan gemaa okoginoon, niisiginoon nishwaaswi (8) ako asho-niizh (12) diba'iganensan (cm)

Endazhiging

Mikigaade bikwaakwaansing, bashkodeng dago zegaakwaasinog

Mawinzon mitigoonsing onji. Miskobagesaanensiwag, okogiwaad bezhig (1) cm akogi gigi-bagishinowaad.

Niibing mawinzon, July aginzod.

Giiyaabi noonde-gikendaman aaniin qe-danaabaji'adwaaban miskominag, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.

Nannyberry (Viburnum lentago)

- Deciduous shrub. Grows up to 6 metres tall
- Opposite branching. Leaves green and coming to a point at the tip

Flowers are small and white in large clusters

The seed is flat with a ridge along the middle

Where to Harvest

Found near rivers and streams.

Often around aspen in woods or bush where it has access to sunlight.

How to Harvest

Harvest by pulling the fruit off the shrub.

Fruit black when ripe.

When to Harvest

Harvest in late summer, August – September.

Uses

Can be eaten raw, and dried for later use.



Aditemin(an)

- Giitwaami-aniibiishiwan. (6) ningodwaaso-diba'iganaatig (m) apiitakide
- Edawaya'ii odikwaniwan. Ashkaabagaawan, jiibwaawan onakwaya'ii
- Waabishki-waabigwaniinsiwanoon wawokoginoon.
- Nabagaa gitigens, bikogin naawaya'ii

Endazhiging

Mikigaade jiigi-ziibi dago ziibiinsing. Memindage azaadiikaang gemaa mekwaakwaang aandi ezhi-zaagaateg.

Mawinzon mitigoonsing onji. Makadewaa apii aditeg.

Mawinzom gizhibiniibing, August ako September aginzowaad.

Pin Cherry (Prunus pensylvanica)

Shrub or small tree, 1-5 m tall, bark reddish brown, peeling in horizontal strips with prominent raised pores

Leaves, oval to lance shaped, 3-10 cm long, gradually taper to long point at tip, rounded at base, edges have small rounded teeth

Flowers, in flat top clusters, white, 6-10 mm

Where to Harvest

Found in forest clearings, hillsides and riverbanks, usually on well drained sites.

How to Harvest

Harvest by pulling the fruit off the shrub. Fruits, bright red cherries, 5-8 mm across, sour.

When to Harvest

Harvest in late summer, August.

Uses

Can be eaten fresh or cooked to make a jelly. Can also be used in making pemmican.

To find out more about other types of uses from the pin cherry medicinal plant offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.





Bawa'iminaan(an)

- Bikwaakwaans gemaa mitigoons, bezhig (1) ako naanan (5) (m) akoonsiinsi, misko-ozaawizi wanagek, bepeshi-gidagizi ezhimamakizid
- Zhaagawaawan gemaa aako-jiibwaawan aniibiishan, niswi (3) ako midaaswi (10) cm akoozi. Onakwaya'ii izhi jiibozi, anaaming waawiyejiiminagizi, gizhibi-aya'ii babikogiinsi
- Oko-nabagi-waabigwaniwi, waabishkizi, 6-10 mm akogi

Endazhiging

Mikigaade bashkodeng, aasimajiw dago jiigibiing, memindage netaa-iskising ningoji.

Mawinzon mitigoonsing onji. Aako-miskoziwag, 5-8 mm iniginiwag, ziiwiskiziwag. Gidaa-ashkamaag gemaa gidaa-giizizwaag jibaashkiminisigeyan.

Mawinzon gizhibiniibing, August aginzod.

Giiyaabi noonde-qikendaman aaniin qe-danaabajiwindwaaban miskominag, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.

Woodland Strawberry (Fragaria vesca)

- Perennial with scaly rhizome and long, slender trialing stolons, leaf stalks and flower stems greenish or very lightly tinged reddish purple, lightly to densely hairy
- Leaves, at stem base, compound, 3 leaflets 10-30 mm long, stalkless strongly toothed,
- Flowers, 3-15 on stem base compound three leaflets, 10-30 cm long, white five petals 8-11 long

Where to Harvest

Found in dry to moist open woods, stream banks, and meadows.

How to Harvest

Harvest by pulling the fruit off the stem. The fruit is red berries. The fruit stems longer than leaves.

When to Harvest

Can be harvested in the summer, July.

Uses

Eaten fresh or preserved for later use. Can be made into a jam.

To find out more about other types of uses from the wild strawberry medicinal plant offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.



Ode'imin(an)

- Endasowaki nitaawigin, ginooginoon ojiibikoon edawaya'ii nikiiginoon, ashkaabaganoon gemaa miskoonaagwadoon, miishaamagadoonsiwanoon
- Wendagigin aniibiishan, okoginoon, nisobagaawan 10-30 mm akoginoon, mitigoonsiwanzinog, mashkawidagaawan
- Waabigwaniwan, niswi (3) ako asho-naanan (15) anaaming ozhiginoon nisobagaawan, 10-30 cm akoginoon, naananobagaawan 8-11 akwaawan

Endazhiging

Mikigaade ningoji baateg gemaa dipaabaaweg nekobaag, jiigi-ziibiinsing dago bashkodeng.

Mawinzon mitigoonsing onji. Miskwaawan ode'eminan. Nawach ginwaawan odikwanensan apiich aniibiishan.

Niibing mawinzom July aginzod.

Giiyaabi noonde-qikendaman aaniin qe-danaabajichigaadegin ode'eminan, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.

Rosehips (Rosa sp.)

- Low growing shrub. Not more than 1.5 metres tall. Stems brown and covered in thorns
- Deciduous. Leaves green and 2 to 5 centimetres long
- Flowers pink and somewhat large, with 5 petals on each flower

Where to Harvest

In open areas, forest and bushes. Common on roadsides.

How to Harvest

Harvest by pulling the fruit from the stem. Careful the stem has many thorns. Fruit red with remnants of flower petals on end opposite the stem.

When to Harvest

Harvest after the first frost.

Uses

Eaten fresh. Be sure to pit (take the seeds out) prior to eating. Can be made into a jelly.

To find out more about other types of uses from rosehip medicinal plant offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.



Oginiimin(an) Oginimingaawanzh

- Zhagakiginoon. Gaawiin awashime 1.5 (m) apiitakidesinoon. Ozaawaa-gaashaakozi
- Giitwaami-aniibiishiwan. Ozhaawashkobagaa (2) ako (5) cm akwaawan
- Miskoonaagwadoon waabigwaniin, nawach igo michaawan, nenaananobagaagin

Endazhiging

Bashkodeng, megwaakwaang dago nekobaang. Jiigikana gaye nitaawiginoon.

Mawinzon mitigoonsing onji. Weweni igo, gaashaagaakozi. Miskojiiminagiziwag, bebangii waabigwaniinsiwanoon anaaming.

Mawinzom ishkwaa nitam mashkawaakwading.

Giiyaabi noonde-qikendaman aaniin qe-danaabajichigaadegin oginiiminensan, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.



Buffalo Berry (Shepherdia canadensis)

- Spreading shrub or small tree, 1-4 m tall, grows from spreading rhizomes, twigs densely covered with rusty brown scales
- Leaves: alternate, oval to egg or lance shaped, 2-6 cm long, densely silvery with tiny shaped hairs on both surfaces
- Flowers, single or in small clusters, yellowish brown

Where to Harvest

Found in open woods, thickets, and riverbanks.

How to Harvest

Harvest by pulling the fruit off of the branch, Fruits, bright red to yellowish, translucent spherical to oval juicy berries, 3-6mm long extremely bitter, soapy to touch.

When to Harvest

Harvest in late summer, August – September.

Uses

Can be made into a jelly and preserved for later use.

This berry also has other uses, offer your tobacco to someone knowledgeable in your community and ask about the plant of the buffalo berry and its medicinal properties.







Mashkode-bizhikiwimin(an)

- Mitigoons gemaa zhazhwegigi, 1-4 (m) da-apiitakizo, zasweginoon ojiibikoon, ozaawi-gidagaawan odikwanensan
- Aniibiishan: zhaagawaawan gemaa jiibwaawan, 2-6 (cm) akwaawan. Miziwe waabi-miishaansiwanoon edawaya'ii
- Waabigwaniin nenishkiginoon gemaa okoginoon, waabizaawaanoon
- Mikigaade nekobaang, bikwaakwaansing dago jiibi-ziibiing

Endazhiging

Mawinzon odikwaning onji, aako-miskwaawan gemaa waabizaawaawan, gegaa zhaabwanaagwadoon, waawiyejiiminagiziwag, 3-6 (mm) akooziwag. Aapiji ziiwiskipogoziwag, zhooshkwajiwag.

Mawinzom giizhibiniibing

Bebakaan gidaa-danaabaji'aawaa mashkode-bizhikiwiminan, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.

Beaked Hazelnut (Corylus cornuta)

- Shrub, 1-3m tall, young twigs, leaves, and bud scales covered in long white hairs, twigs hairless after first season
- Leaves, elliptic to egg shaped, 4-10 cm long. Rounded to heart shaped at the base, sharp pointed at the tip, edges are coarsely double toothed

Where to Harvest

Found in moist but well drained sites in thickets or woods.

How to Harvest

Pull the husks off the plant. Fruit is thin shelled, spherical, edible nuts enclosed in long tubular husks, husk light green, covered with stiff, prickly hairs, narrowly lobed at tip, in clusters of 2 or 3 at ends of branches.

When to Harvest

Harvest in late summer.

Uses

The husks can be cleaned so you can roast the nuts with seasonings.

You also can also offer your tobacco to someone knowledgeable in your community and ask about the plant of the hazelnut and its medicinal properties.





Jiibwebagaan(ag)

- Mitigoons, 1-3 (m) akoozi, oshki-odikwaniwan, aniibiishiwan, oshkibagaag waabishki-miishaansiwan, nitam nitaawigig, gaawiin miishaanziwanzinoon
- Zhaagawaawan aniibiishan, 4-10 (cm) akwaawan. Nawach igo waawiyeyaa anaaming, jibwaawan onakwaya'ii, neniizhogidagaawan edawaya'ii
- Ningoji dipaabaaweg netaa-iskising bikwaakwaansing gemaa megwaakwaang

Endazhiging

Gizhibiniibing mawinzom, gibishigibinaag. Bibagiziwag, waawiyeziwag, amogaazowag, ginoo-gibogiwag, waabishki-ashkaabagizi, miishaansiwag, ani-jiiboziwag onakwaya'ii. Neniizhigiwag (2) gemaa nenisigiwag (3) onakwaatigoonsing. Apii bishagibinadwaa, gidaa-abwe bagaanag. Gidaa-zhiiwitaaganiwaag gaye.

Giiyaabi noonde-aikendaman aaniin ge-danaabajiwindiban bagaan, miizh asemaa gichi-anishinaabe gemaa awiya gekendana gechiwaag aki.

Sage (Aretemisia frigida)

- A silvery gray aromatic perennial with a woody crown and root, growing up to 50cm tall
- Spreading to form a mat of, with fuzzy stems, basal leaves that are divided into narrow segments

Where to Harvest

Found in meadows and pastures in the parklands.

How to Harvest

Harvest by snapping the stem near the earth. Hang bunches of sage to dry.

When to Harvest

Harvest in summer, July or August.

Uses

Commonly, people use this to smudge.

To find out more about other types of uses from the sage medicinal plant offer your tobacco and speak with an elder or someone knowledgeable about the land in your community.



Mashkodebizhikii-wiingwashkosigan(oon)

- Nawach igo waabanaagwad, minomaagwad, mitigowe-ayaamagad, ojiibikoomagad, 50 (cm) apiitakide
- Zaswe-miishaansiwanoon aniibiishensan, bebikish agaasideyaawan

Endazhiging

Mikigaade mashkodeng dago bashkodeng.

Odaapinigaadeg, wendaging gibookobidoon azhishkiing. Agoodoon jibaasaman.

Gidaa-moozhiginaan niibing July gemaa August aginzowaad.

Giiyaabi noonde-qikendaman aaniin endanaabadak mashkodewiingashk, miizh asemaa gichi-anishinaabe gemaa awiya gekendang gechiwaag aki.

Sweet Flag (Acorus americanus)

Aquatic perennial herb growing from an aromatic rhizome, with long narrow sword shaped leaves, with a leaf like flower stalk that produces a dense cylindrical spike of tiny yellow flowers

Where to Harvest

Found in scattered locations in swamps, along water ways, and in the boreal forest.

How to Harvest

Harvest by following the stems to the root and digging out the root. The root then can be cleaned and dried.

When to Harvest

Harvest in the early fall, September.

Uses

Can be eaten in small pieces after it is dried and also used to make a tea.

This plant also has other uses, offer your tobacco to someone knowledgeable in your community and ask about the sweet flag plant and its medicinal properties.



Wiike

Niibing nitaa-nitaawigin ojiibik, jiibwaawan ginwaagin aniibiishan, daabishkoo waabigwan ojiibiik, agaasinging waabizaawaagin waabigwaniinsan gizhibaaginoon

Endazhiging

Niibing nitaa-nitaawigin ojiibik, jiibwaawan ginwaagin aniibiishan, daabishkoo waabigwan ojiibiik, agaasinging waabizaawaagin waabigwaniinsan gizhibaaginoon.

Mashkiigong mikigaade, jiigibiig dago megwaakwaang.

Gidaa-odaapinaan nisidawinaman, mii ezhi-gidakiibidooyan ojiibiik. Gibiinitoon ojiibiik, gidaa-baasaan.

Wiiba dagwaagig odaapinan, September aginzod.

Giishin noonde-qikendaman aaniin endanaabadak owe ojiibiik, miizh asemaa qichi-anishinaabe qemaa awiya qekendang qechiwaaq aki jiwiindamaag aaniin ji-izhichigaadaman.



Muskeg/Labrador Tea

(Ledum groenlandicum)

- A stout shrub with alternate evergreen leaves. Edges roll under and have rusty hair underneath
- Flowers are white, with small five toothed sepal tubes and five pedals, fruit is small and fuzzy

Where to Harvest

Found in muskeg, bogs and wet coniferous woods.

How to Harvest

Harvest by pulling the leaves from the stems. Leaves can be dried for later use.

When to Harvest

Labrador tea can be harvested in every season.

Uses

Commonly used to make a tea.

This plant also has other uses, offer your tobacco to someone knowledgeable in your community and ask about the plant and its medicinal properties.



Omashkiigobag(oon)

- Okogin netaawigig, bebakaan ayashkaabigwaniiwan. Anaaming izhigin ozaawaanikwemagad anaaming
- Waabishki-waabigwaniiwan, naanwaabidewan dago naanibagiwaniinsiwan, agaansin miishaanziwan gaye

Endazhiging

Mikigaade mashkiig, dootoobanong gemaa zhaabobiig zhingobiikaang.

Gidaa-odaapinaanan aniibiishan odikoonsing onji. Gidaa-baasaanan aniibiishan naagach onji.

Bigo apii gidaa-moozhaginaanan omashkiibagoon.

Bebakaan gaye inaabadad owe mashkiibag, miizh asemaa gichianishinaabe gemaa awiya gekendang gechiwaag aki, giga-wiindamaag aaniin ge-izhichigaadaman.

Sweet Grass (Hierochloe odorata)

A perennial sweet smelling grass growing up to 60 cm tall. The leaf sheaths are smooth, blades are lance shaped and up to 6mm wide

Where to Harvest

Found in meadows and wet ground.

How to Harvest

Harvest by not pulling the root out, but by snapping the stem near the earth. The strands can be braided and dried.

When to Harvest

Harvest in the later summer, July or August.

Uses

Commonly, people use this to smudge.

This plant has other uses, offer your tobacco to someone knowledgeable in your community and ask about the plant of the sweet grass and its medicinal properties.



Wiingwashkosigan(oon)

Endasowaki nitaawigin mashkosiw. 60 (cm) apiitakide. Zhooshkwaawan aniibiishan, nawach igo jiibwaawan 6 (mm) inigokwaawan

Endazhiging

Mashkodeng mikigaade gemaa zhaabobiikamagaang.

Gidaa-odaapinaan, gego gidakiiken, bookonan wendaging jiigikamig. Gidaa-apikaadaanan ge-izhi-baasaman.

Gidaa-moozhaginaan ani-gizhibiniibing, July gemaa August aginzowaad.

Bebakaan danaabadadoon wiingashkoon, miizh asemaa gichianishinaabe gemaa awiya gekendang gechiwaaq aki, qiqa-wiindamaaq aaniin ji-izhichiqaadaman.

Preparing to harvest

It is important to let someone know where you will be going to gathering. Prepare for yourself and whomever you are going to go gathering with. When you leave your gathering areas, make sure you don't leave anything behind, leave it as you found it.

What to Bring...

- Tobacco, to make your offering.
- **Back pack,** to put your snack, water and miscellaneous items.
- **Bucket,** to put your fruit in.
- **Blanket,** to wrap your medicines in.
- **Scissors,** to clip your harvest, so roots stay in the ground.
- String, to tie medicine together.
- Cloth, to hold medicine together.
- Knife, to cut string or cloth.
- Boots, to keep your feet dry.
- Jacket, to keep yourself dry.









"Standing Medicine People: A guide to common medicines found in south central Manitoba" was made in partnership with Long Plain First Nation, Swan Lake First Nation, Rolling River First Nation, and Food Matters Manitoba.

Learn more at: foodmattersmanitoba.ca

Created by Food Matters Manitoba © 2015. Many thanks to MFNERC for printing and to the University of Manitoba Herbarium (WIN) for use of their photos. Chokecherries, Wild Raspberry, Pin Cherry, Woodland Strawberry, Rosehips and Sweet Grass © iStock photos, p. 8, 10, 14, 16, 18, 30, respectively. Cover illustration by Moi & Toi Design. Guidebook content based on Aboriginal Plant Use in Canada's Northwest Boreal Forest (UBC Press: 2000) and Plants of the Western Boreal Forest & Aspen Parkland (Lone Pine Publishing: 1995).